

# The State of the World's Antibiotics in 2018


Ella Pringle Lecture  
Royal College of Physicians of Edinburgh

Ramanan Laxminarayan  
Twitter @CDDEP



- I. Drug resistance is rising worldwide and threatens gains made in reducing the burden of infectious diseases

## First reported cases of bacterial resistance against key antibiotics



Data source: Antibiotic Resistance Threats in the United States, 2013, US Centers for Disease Control and Prevention (CDC)



It is not difficult to make microbes resistant to penicillin in the laboratory by exposing them to concentrations not sufficient to kill them, and the same thing has occasionally happened in the body...

*Alexander Fleming, 1945*

## Antibiotic resistance is ancient


Vanessa M. D'Costa<sup>1,2\*</sup>, Christine E. King<sup>3,4\*</sup>, Lindsay Kalan<sup>1,2</sup>, Mariya Morar<sup>1,2</sup>, Wilson W. L. Sung<sup>4</sup>, Garsten Schwarz<sup>3</sup>, Duane Froese<sup>5</sup>, Grant Zazula<sup>6</sup>, Fabrice Calmebs<sup>5</sup>, Regis Debruyne<sup>7</sup>, G. Brian Golding<sup>4</sup>, Hendrik N. Poinar<sup>1,2,4</sup> & Gerard D. Wright<sup>1,2</sup>

The discovery of antibiotics more than 70 years ago initiated a period of drug innovation and implementation in human and animal health and agriculture. These discoveries were tempered in all cases by the emergence of resistant microbes<sup>1,2</sup>. This history has been interpreted to mean that antibiotic resistance in pathogenic bacteria is a modern phenomenon; this view is reinforced by the fact that collections of microbes that predate the antibiotic era are highly susceptible to antibiotics<sup>3</sup>. Here we report targeted metagenomic analyses of rigorously authenticated ancient DNA from 30,000-year-old Beringian permafrost sediments and the identification of a highly diverse collection of genes encoding resistance to  $\beta$ -lactam, tetracycline and glycopeptide antibiotics. Structure and function studies on the complete vancomycin resistance element VanA confirmed its similarity to modern variants. These results show conclusively that antibiotic resistance is a natural phenomenon that predates the modern selective pressure of clinical antibiotic use.

Recent studies of modern environmental and human commensal microbial genomes have a much larger concentration of antibiotic resistance genes than has been previously recognized<sup>4-6</sup>. In addition,

with high concentrations of *Escherichia coli* harbouring the *gfp* (green fluorescent protein) gene from *Aequorea victoria* (Supplementary Information).

After fracturing of the samples (Supplementary Fig. 3), total DNA was extracted from a series of five subsamples taken along the radius of each core (Supplementary Information). Quantitative polymerase



**MONEY WEEK**

The w  
super

**WHITWATER**  
**News**

**ANT**

THE EP

# COSMOPOLITAN

November 1995

At Last!  
Something  
Pleasurable  
That's  
Good  
for You.  
**The Health  
Benefits  
of Sex**

Cosmo's  
Update on  
Antibiotics.  
What's Okay  
and What's  
Dangerous

The  
Heart-  
Pounding  
Bawdiness  
of  
**Brad  
Pitt,**  
Who  
Couldn't  
Care  
Less

**Why  
Marry  
Instead of  
Just  
Fooling  
Around?**

**Makeup Tricks**

\$2.95



**ESTING** MUTUAL FUNDS:  
1st QUARTER STARS  
**Week**

\$3.95

gainst  
e  
**BES**  
g back against  
tious disease.

DR. DAVID SHLAES  
OF WYETH-AYERST

## Carbapenem and 3rd. gen. cephalosporin resistance among *K. pneumoniae* highest along the East Coast, but present in all regions of the country

### Carbapenem




### 3rd Gen. Cephalosporins




Note: Data for 2010 available through July.

Data source: Braykov NB, Eber MR, Klein EY, Morgan DJ, Laxminarayan R. Trends in Resistance to Carbapenems and Third- Generation Cephalosporins among Clinical Isolates of *Klebsiella pneumoniae* in the United States, 1999-2010. *Infect Control and Hospital Epidemiology*. 2013; 34(3)






CRE rates in children grew between 2000 and 2012



Logan et al, EID, 2015





**Trends in antimicrobial resistance in bloodstream infection isolates at a large urban hospital in Malawi**

Musicha et al, *Lancet Inf Dis* 2017

# Carbapenem-resistant *Acinetobacter baumannii*

Year: **1999**

Percentage resistant:

 < 10%


 10 - 20%

 20 - 40%


 40 - 60%

 > 60%

 Inadequate data




## Percentage of *Staphylococcus aureus* that are methicillin resistant (MRSA), by country (most recent year, 2011-14)



Source: CDDEP 2015, WHO 2014 and PAHO, forthcoming

Where available, data from hospital-associated MRSA and invasive isolates have been used. In their absence, data from community-associated MRSA or all specimen sources are included. Only countries that reported data for at least 30 isolates are shown. Depending on the country, resistance to one or more of the following drugs were used to test for MRSA: Oxacillin, cefoxitin, flucloxacillin, cloxacillin, dicloxacillin, and methicillin. Intermediate-resistant isolates are included as resistant in some calculations, as in the original data source.

## Percentage of extended-spectrum beta-lactamase producing *Escherichia coli*\*, by country (most recent year, 2011-2014)




Source: CDDEP 2015, WHO 2014 and PAHO, forthcoming

Where available, data from invasive isolates have been used. In their absence, data from all specimen sources are included. Only countries that reported data for at least 30 isolates are shown. Depending on the country, resistance to one or more of the following drugs were used: Ceftazidime, ceftriaxone and cefotaxime. Intermediate-resistant isolates are included as resistant in some calculations, as in the original data source.


\*Indicated by third-generation cephalosporin resistance

## Percentage of carbapenem-resistant *Klebsiella pneumoniae*, by country (most recent year, 2011-2014)



Source: CDDEP 2015, WHO 2014 and PAHO, forthcoming

Where available, data from invasive isolates have been used. In their absence, data from all specimen sources are included. Only countries that reported data for at least 30 isolates are shown. Depending on the country, resistance to one or more of the following drugs were used: imipenem, meropenem, ertapenem and doripenem. Intermediate-resistant isolates are included as resistant in some calculations, as in the original data source.



Woerther, Clin Microbiol Rev. 2013

RECENT ARTICLES

A Dow Jones for Drug Resistance

by Martin Enserink on 5 November 2010, 3:14 PM | Permanent Link | 1 Comments


MORE EVIDENCE FOR HIDDEN PARTICLES?

A DOW JONES FOR DRUG RESISTANCE

PODCAST: LIFELIKE

Email Print Facebook Twitter YouTube LinkedIn StumbleUpon More

PREVIOUS ARTICLE NEXT ARTICLE




Bullish or bearish? The new index could track resistance in Salmonella and many other bacterial pathogens.

Credit: B.Thomason/CDC

## The drugs work less

Effectiveness of antibiotics\* in selected countries  
0=best, 100=worst



Source: Centre for Disease Dynamics, Economics & Policy

Economist.com

\*By index of six bacteria's resistance to six classes of drug

<sup>1</sup>Except US (2012), Poland (2013) and Thailand (2013)



# Staphylococcus aureus resistance to oxacillin (%MRSA)







FIGURE 1. Percentage of *Escherichia coli* and *Klebsiella pneumoniae* isolates from selected health care-associated infections\* with the extended-spectrum- $\beta$ -lactamase (ESBL) phenotype reported as nonsusceptible to extended-spectrum cephalosporins† — National Healthcare Safety Network, United States, 2006–2015



\* Central line-associated bloodstream infections and catheter-associated urinary tract infections.

† Nonsusceptible to at least one extended-spectrum cephalosporin.

FIGURE 2. Percentage of *Escherichia coli* and *Klebsiella pneumoniae* isolates from selected health care–associated infections\* reported as resistant to a carbapenem — National Healthcare Safety Network, United States, 2006–2015



\* Central line–associated bloodstream infections and catheter-associated urinary tract infections.




Figure 1. Carbapenem- and cephalosporin-resistant *Acinetobacter baumannii* in the United States: 1999–2012.

Logan et al *J Ped Inf Dis Soc*, 2018.

# Are we overstating the problem of resistance because of measurement bias?



Clinical Infectious Diseases

## BRIEF REPORT

### Is Antimicrobial Resistance a Bigger Problem in Tertiary Care Hospitals Than in Small Community Hospitals in the United States?

Samanth Gandra,<sup>1\*</sup> Anna Trett,<sup>1,4</sup> Eili Y. Klein,<sup>2,3</sup> and Ramanan Laxminarayan<sup>2,4</sup>

<sup>1</sup>Center for Disease Dynamics, Economics & Policy, New Delhi, India; <sup>2</sup>Center for Disease Dynamics, Economics & Policy, Washington, DC; <sup>3</sup>Johns Hopkins University, Department of Emergency Medicine, Baltimore, Maryland, and <sup>4</sup>Princeton Environmental Institute, New Jersey




-- Outpatient, community

— Inpatient, tertiary


-- Outpatient, tertiary

— Inpatient, community

## Spread of New Delhi metallo beta-lactamase: first detection, by country




Source: Johnson and Woodford 2013 (adapted)



**Antimicrobial susceptibility pattern of 361 *Streptococcus pneumoniae* isolates**

Manoharan LID, 2017


## Numbers of unique $\beta$ -lactamase enzymes identified since introduction of first $\beta$ -lactam antibiotics



Davies and Davies, Microbiol. Mol. Biol. Rev. 2010.



# Mortality outcomes are worse in neonates with resistant infections



Kayange M, Kamugisha E, Mwizamholya DL, Jeremiah S, Mshana SE. 2010. Predictors of positive blood culture and deaths among neonates with suspected neonatal sepsis in a tertiary hospital, Mwanza- Tanzania. BMC Pediatrics 10: 39.

	Number of resistant isolates	CFR in culture-positive sepsis due to resistant pathogens	CFR in culture-positive sepsis due to sensitive pathogens
<b>Gram negative</b>			
<i>Acinetobacter</i> spp (n=222)			
ES cephalosporins	85/222 (38%)	59/85 (69%)	71/137 (52%)
Carbapenems	174/222 (78%)	106/174 (61%)	24/48 (50%)
MDR	181/222 (82%)	112/181 (62%)	18/41 (44%)
<i>Klebsiella</i> spp (n=169)			
ES cephalosporins	105/169 (62%)	57/104 (55%)	38/65 (58%)
Carbapenems	59/169 (35%)	36/59 (61%)	59/110 (54%)
MDR	91/169 (54%)	52/91 (57%)	43/78 (55%)
<i>Escherichia coli</i> (n=137)			
ES cephalosporins	65/137 (47%)	40/64 (63%)	43/73 (59%)
Carbapenems	21/137 (15%)	12/21 (57%)	71/116 (61%)
MDR	52/137 (38%)	30/52 (58%)	53/85 (62%)
<i>Pseudomonas</i> spp (n=68)			
ES cephalosporins	32/68 (47%)	29/32 (91%)	24/36 (67%)
Carbapenems	21/68 (31%)	19/21 (90%)	34/47 (72%)
MDR	13/68 (19%)	11/13 (85%)	42/55 (76%)

DeNIS Study, Lancet ID, 2016





Figure: Case fatality rates from the DeNIS study (unshaded) compared with earlier studies (in solid colours)



**Figure 2: Estimated neonatal sepsis deaths caused by bacteria resistant to first-line antibiotics in five high-burden countries**

Laxminarayan et al Lancet, 2015

"All the News  
That's Fit to Print"

# The New York Times

Late Edition

Today, a chilly day, sunshine mixing with some clouds, high 41. Tonight, mostly cloudy, low 34. Tomorrow, mostly cloudy, chilly, high 41. Weather map is on Page B4.

VOL. CLXIV ... No. 56,705 +

© 2014 The New York Times

NEW YORK, THURSDAY, DECEMBER 4, 2014

\$2.50



Protesters at Grand Central Terminal on Wednesday after a grand jury decided not to indict a police officer in Eric Garner's death.

## NEW YORK OFFICER FACING NO CHARGES IN CHOKEHOLD CASE

### Grand Jury's Decision in Fatal Encounter Draws Protests — U.S. to Investigate

By J. DAVID GOODMAN and AL BAKER

A Staten Island grand jury on Wednesday ended the criminal case against a white New York police officer whose chokehold on an unarmed black man led to the man's death, a decision that drew condemnation from elected officials and touched off a wave of protests.

The fatal encounter in July was captured on video and seen around the world. But after viewing the footage and hearing from witnesses, including the officer who used the chokehold, the jurors deliberated for less than a day before deciding that there was not enough evidence to go forward with charges against the officer, Daniel Pantaleo, 29, in the death of the man, Eric Garner, 43.

Officer Pantaleo, who has been on the force for eight years, appeared before the grand jury on Nov. 21, testifying that he did not intend to choke Mr. Garner, who was being arrested for allegedly selling loose cigarettes. He described the maneuver as a take-down move, adding that he never thought Mr. Garner was in mortal danger. [Page A28.]

The decision came barely a week after a grand jury found no criminality in the actions of another white police officer, Darren Wilson, who shot and killed Michael Brown, an unarmed 18-year-old black man in Ferguson, Mo.

After the news from Staten Island, a wave of elected officials renewed calls for Justice Department intervention, saying the

grand jury's finding proved that justice could be found only in the federal courts. By the evening, the department announced it would open a civil rights inquiry.

On the streets of the city, from Times Square to Times Square, many expressed their outrage with some of the last words Mr. Garner uttered before being wrestled to the ground: "This stops today," people chanted. "I can't breathe," others shouted.

While hundreds of angry but generally peaceful demonstrators took to the streets in Manhattan as well as in Washington and other cities, the police in New York reported relatively few arrests, a stark contrast to the riots that unfolded in Ferguson in the hours after the grand jury decision was announced in the

Continued on Page A28



Mr. Garner, in an undated family photo, died at age 43.

## U.S. and Iran Both Attack ISIS, But Try Not to Look Like Allies

By TIM ARANGO and THOMAS ERDBRINK

BAGHDAD — Iranian fighter jets struck extremist targets in Iraq recently, Iranian and American officials have confirmed, in the latest display of Tehran's new willingness to conduct military operations openly on foreign battlefields rather than covertly and through proxies.

The shift stems in part from Iran's deepening military role in Iraq in the war against the Sunni extremists of the Islamic State. But it also reflects a profound

shift in a buffer zone that extends 25 miles into Iraq.

The new military approach highlights an unusual confluence of interests in both Iraq and Syria, where Tehran and Washington find themselves fighting the same enemy in an increasingly public fashion. While there is no direct coordination between Iran and the United States, there is a de facto nonaggression pact that neither side is eager to acknowledge.

## 'Superbugs' Kill India's Babies And Pose an Overseas Threat

By GARDNER HARRIS


AMRAYATI, India — A deadly epidemic that could have global implications is quietly sweeping India, and among its many victims are tens of thousands of newborns dying because antimicrobial drugs no longer work.

These infants are born with bacterial infections that are resistant to most known antibiotics, and more than 58,000 died last year as a result, a recent study found. While that is still a fraction of the nearly 800,000 newborns

world, and this will require treating an increasing number of newborns who have sepsis and pneumonia," said Dr. Vinod Paul, chief of pediatrics at the All India Institute of Medical Sciences and the leader of the study. "But if resistant infections keep growing, that progress could slow, stop or even reverse itself. And that would be a disaster for not only India but the entire world."


It starts to neonatal intensive care wards in five Indian states,

Absolute risk reduction (ARR) of infection with antibiotic prophylaxis in common surgical procedures and blood cancer chemotherapy in the USA



Teillant et al, *Lancet Infect Dis*, 2015

## Number of additional infections per year in the USA under a 30% decreased efficacy of antibiotic prophylaxis



Teillant et al, *Lancet Infect Dis*, 2015

# UTIs turn deadly with reckless antibiotic use

Bangalore, India  
@timesofindia

New Delhi: Indiscriminate use of antibiotics in India is making common infections like urinary tract infection (UTI) deadly. Urologists say one out of every 10 UTI patients are now requiring life-line drugs and, in some cases, even that is failing to help.

"We are forced to use old reliable antibiotics with known toxicity to kidneys, for example Chloramphenicol, to save patients with multi-drug resistant UTI," Dr Anoop Kumar, professor and head of the urology department at Safford Hospital, said.

UTI is one of the most common bacterial infections that affect one or more parts of the urinary system. It is more common in women than men.

Kumar said that if steps are not taken to strictly regulate and monitor prescription and sale of antibiotics immediately the bacteria triggering UTI will turn resistant to even the remaining drugs. "People may die of UTI," the Safford Hospital doctor added.

Urologists at the Gupta, Ravi and BLK Super-specialty hospitals have also confirmed the trend of rising drug resistance in UTI cases.

"Four to five patients die of kidney failure due to UTI infection each year. It is either caused by the infection itself or heavy dosage of antibiotics administered to treat them," Dr Anoop Kumar, the

## POINT OF NO RETURN

Antibiotic resistance is a global health crisis.

It's the resistance of an antibiotic drug that was originally effective in treating infections caused by bacteria.

### CAUSES

- Over-prescription of antibiotics
- Overuse of antibiotics in livestock and fish farming
- Poor infection control in hospitals & clinics

head of urology and kidney transplant department at All India Institute of Medical Sciences (AIIMS), said.

According to Dr Anoop Kumar, Professor, senior urology consultant at the Sir Ganga Ram Hospital, most UTIs are caused by gram-negative bacteria such as E. coli.

"UTI about 10 years ago, most UTIs were treatable with oral antibiotics. It was less toxic and easy on the pocket, but due to drug resistance many patients have to be admitted and they are put on intravenous drugs that are five to six times costlier," he said.


Women are at higher risk of UTI due to anatomy of the female system, which is 1.5 inches long compared to 8 inches in men. Pregnancy or recent sexual activity and other causes are other common causes that lead to UTI after stopping it, when they begin to develop possible problems.



# Surgical site infections


- There are 92 million surgeries in low-income countries each year
- 5.5 million surgical site infections or SSIs (6 per 100 procedures) – about a third of all healthcare associated infections
- SSIs are the leading cause of infection in settings with limited resources
- Mortality rate from SSI Rates of mortality from surgical site infections are 3% in the US and between 8 and 20% in low-income countries
- Between 400,000 and a million deaths from SSIs each year with an increasing number caused by resistant pathogens.

**Zero to zero in 100 years:  
Available resistance-free antimicrobial classes for *Neisseria gonorrhoeae***



II. Rising incomes and increasing access to antibiotics are saving lives (although lack of access still kills more people than antibiotic resistance) but are not a good substitute for public health

Bacterial diseases are still major killers in developing countries because of lack of access to antibiotics




O'Brien et al, Lancet 2009

# What are we asking of antibiotics?

FIGURE 1.1

Crude infectious disease mortality rate in the United States, 1900–1996




Source: Adapted from Armstrong, Conn et al. (1999).

# Substitute for immunization, infection control and water/sanitation

FIGURE 1.1

Crude infectious disease mortality rate in the United States, 1900–1996




Source: Adapted from Armstrong, Conn et al. (1999).

# Substitute for immunization, infection control and water/sanitation

FIGURE 1.1


Crude infectious disease mortality rate in the United States, 1900–1996



Source: Adapted from Armstrong, Conn et al. (1999).

# Vaccines can be effective


Invasive disease caused by Pneumococci in children under two declined in the US post pneumo vaccination



Kyaw MH et al. N Engl J Med 2006;354:1455-1463.




## Effect of PCV7 introduction in 2000 on antibiotic prescriptions and ambulatory care visits




Zhou et al, Pediatrics 2008

## Effect of PCV7 introduction in 2000 on antibiotic prescriptions and ambulatory care visits

Antibiotic prescriptions attributable to acute otitis media decreased from 1244 to 722 prescriptions per 1000 person-years – a 41.9% reduction.



# Antibiotic consumption is increasing in developing countries...



Source: Based on data obtained under license from IMS Health MIDAS™ (January 2005-December 2010); IMS Health Incorporated. All Rights Reserved.

---

# Global antibiotic consumption 2000 to 2010: an analysis of national pharmaceutical sales data



Thomas P Van Boeckel, Sumanth Gandra, Ashvin Ashok, Quentin Caudron, Bryan T Grenfell, Simon A Levin, Ramanan Laxminarayan

## Summary

**Background** Antibiotic drug consumption is a major driver of antibiotic resistance. Variations in antibiotic resistance across countries are attributable, in part, to different volumes and patterns for antibiotic consumption. We aimed to assess variations in consumption to assist monitoring of the rise of resistance and development of rational-use policies and to provide a baseline for future assessment.

**Lancet Infect Dis 2014**  
Published Online  
July 30, 2014  
<http://dx.doi.org/10.1016/j.slanid.2014.07.007>




# Global increase and geographic convergence in antibiotic consumption between 2000 and 2015


Eili Y. Klein<sup>a,b,c,1</sup>, Thomas P. Van Boeckel<sup>d</sup>, Elena M. Martinez<sup>a</sup>, Suraj Pant<sup>a</sup>, Sumanth Gandra<sup>a</sup>, Simon A. Levin<sup>e,f,g,1</sup>, Herman Goossens<sup>h</sup>, and Ramanan Laxminarayan<sup>a,f,i</sup>

<sup>a</sup>Center for Disease Dynamics, Economics & Policy, Washington, DC 20005; <sup>b</sup>Department of Emergency Medicine, Johns Hopkins School of Medicine, Baltimore, MD 21209; <sup>c</sup>Department of Epidemiology, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD 21205; <sup>d</sup>Institute of Integrative Biology, ETH Zürich, CH-8006 Zürich, Switzerland; <sup>e</sup>Department of Ecology and Evolutionary Biology, Princeton University, Princeton, NJ 08544; <sup>f</sup>Princeton Environmental Institute, Princeton University, Princeton, NJ 08544; <sup>g</sup>Beijer Institute of Ecological Economics, SE-104 05 Stockholm, Sweden; <sup>h</sup>Laboratory of Medical Microbiology, Vaccine & Infectious Diseases Institute, University of Antwerp, 2610 Antwerp, Belgium; and <sup>i</sup>Department of Global Health, University of Washington, Seattle, WA 98104

## Global antibiotic consumption by country: 2000–2015.




**Global antibiotic consumption by country income classification: 2000–2015.**



Eili Y. Klein et al. PNAS doi:10.1073/pnas.1717295115



**Antibiotic consumption rate for HICs, LMICs-UM, and LMICs-LM of the four most-consumed therapeutic classes of antibiotics in DDDs per 1,000 inhabitants per day.**




Eili Y. Klein et al. PNAS doi:10.1073/pnas.1717295115

©2018 by National Academy of Sciences

PNAS

**Antibiotic consumption rate for HICs, LMICs-UM, and LMICs-LM of new and last-resort antibiotics in DDDs per 1,000 inhabitants per day.**




Eili Y. Klein et al. PNAS doi:10.1073/pnas.1717295115

PNAS




# Countries with highest increase in consumption

RANK COUNTRY




Rank according to volume increase; Source: Proceedings of the National Academy of Sciences

## Antibiotic use per capita by income in selected countries, 2010



Source: Van Boeckel et al. 2014 (based on IMS MIDAS) and World Bank 2015


## Carbapenem retail sales in selected countries, 2005–2010 (per 1,000 population)



Source: Laxminarayan et al. 2013 (based on IMS MIDAS)

\*An IMS grouping of Benin, Burkina Faso, Cameroon, Côte d'Ivoire, Gabon, Guinea, Mali, Republic of the Congo, Senegal, and Togo


**Fig 3. Commonly used antibiotic consumption in different regions of China by class in 2011 and 2015.**



Wushouer H, Tian Y, Guan XD, Han S, Shi LW (2017) Trends and patterns of antibiotic consumption in China's tertiary hospitals: Based on a 5 year surveillance with sales records, 2011-2015. PLOS ONE 12(12): e0190314. <https://doi.org/10.1371/journal.pone.0190314>


<http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0190314>

Faropenem consumption has increased by 154% since it was approved for use in India in 2010



Gandra et al, Clin Inf Dis, 2016

# Carbapenem consumption in the hospital sector in selected European countries, 1997–2013



# Non-prescription use of antimicrobials is common




Figure 2: Frequency of non-prescription use of antimicrobials in the general population based on published works. In small areas, countries with similar frequency of non-prescription antimicrobial use have been grouped.





**Table 1. Workforce of Doctors and Nurses According to Country or Region in 2010.\***

<b>Country or Region</b>	<b>Population <i>in millions</i></b>	<b>Doctors <i>in thousands</i></b>	<b>Nurses</b>	<b>Doctors and Nurses/ 1000 Population</b>	<b>Nurse-to-Doctor Ratio</b>
<b>Country</b>					
China	1338	1915	1,864	2.8	0.97
India	1225	768	1,179	1.6	1.54
United States	309	756	3,064	12.3	4.05
Brazil	195	338	1,278	8.3	3.78
United Kingdom	62	166	626	12.7	3.77
South Africa	50	37	198	4.7	5.30
<b>Region</b>					
Americas	937	1974	4,947	7.4	2.5
Europe	899	2744	5,870	9.6	2.1
Middle East and North Africa	590	654	894	2.6	1.4
Southeast Asia	1795	997	1,810	1.6	1.8
Sub-Saharan Africa	847	150	778	1.1	5.2
Western Pacific	1821	2696	3,814	3.6	1.4
World	6888	9216	18,114	4.0	2.0


\* A doctor or nurse is defined as a person with the appropriate qualifications recognized in his or her own country. In this table, the nurse workforce includes nurses and midwives. Data are from the World Health Organization.<sup>9</sup>

**Table 1. Workforce of Doctors and Nurses According to Country or Region in 2010.\***

Country or Region	Population <i>in millions</i>	Doctors <i>in thousands</i>	Nurses	Doctors and Nurses/ 1000 Population	Nurse-to-Doctor Ratio
<b>Country</b>					
China	1338	1915	1,864	2.8	0.97
India	1225	768	1,179	1.6	1.54
United States	309	756	3,064	12.3	4.05
Brazil	195	338	1,278	8.3	3.78
United Kingdom	62	166	626	12.7	3.77
South Africa	50	37	198	4.7	5.30
<b>Region</b>					
Americas	937	1974	4,947	7.4	2.5
Europe	899	2744	5,870	9.6	2.1
Middle East and North Africa	590	654	894	2.6	1.4
Southeast Asia	1795	997	1,810	1.6	1.8
Sub-Saharan Africa	847	150	778	1.1	5.2
Western Pacific	1821	2696	3,814	3.6	1.4
World	6888	9216	18,114	4.0	2.0

\* A doctor or nurse is defined as a person with the appropriate qualifications recognized in his or her own country. In this table, the nurse workforce includes nurses and midwives. Data are from the World Health Organization.<sup>9</sup>

# The flu season is a key driver of antibiotic consumption



Van Boeckel et al, Lancet Inf Dis, 2014

# Influenza in the United States is nearly perfectly predicted by antibiotic sales data





FIGURE 1. Observed and fitted antibiotics series from 2000 to 2007. The solid line represents the actually observed antibiotics series; the dashed line represents the fitted antibiotics series from the time series regression model that uses influenza-like illness as an explanatory series.

# Hospital use of carbapenems is rapidly growing



Source: Based on data obtained under license from IMS Health MIDAS™ (January 1999-December 2010); IMS Health Incorporated. All Rights Reserved.

# Health insurance increases prescribing

Table 4. Use of oral, injected, and all antibiotics per person per year by level of family income and insurance plan

Antibiotic use and income tertile*	Free plan (N = 1935)		Cost-sharing plans (N = 3830)		Ratio of free to cost-sharing (95% confidence interval)†
	Number of antibiotics	Number per person	Number of antibiotics	Number per person	
<b>Oral antibiotics</b>					
Upper one-third	548	0.94	723	0.58	1.63 (1.55, 1.72)
Middle one-third	577	0.93	669	0.57	1.62 (1.53, 1.71)
Lower one-third	442	0.72	386	0.33	2.17 (1.97, 2.39)
All incomes	1670	0.85	1825	0.48	1.79 (1.72, 1.86)
<b>Injected antibiotics</b>					
Upper one-third	45	0.08	89	0.07	1.09 (0.77, 1.54)
Middle one-third	69	0.11	75	0.06	1.73 (1.27, 2.36)
Lower one-third	38	0.06	45	0.04	1.60 (1.05, 2.44)
All incomes	187	0.10	221	0.06	1.67 (1.39, 2.01)
<b>All antibiotics</b>					
Upper one-third	593	1.02	812	0.65	1.57 (1.51, 1.63)
Middle one-third	646	1.04	744	0.64	1.63 (1.57, 1.70)
Lower one-third	480	0.78	431	0.37	2.11 (1.94, 2.30)
All incomes	1857	0.96	2046	0.53	1.80 (1.75, 1.86)

\*Numbers shown for income tertiles do not sum to totals because income was unknown for 138 claims on the free plan and 59 on the cost-sharing plans.

†Taylor's series 95% confidence intervals [12]; ratio and confidence intervals calculated using 8 significant digits.

## Association between health insurance and antibiotics prescribing in four counties in rural China


Hengjin Dong <sup>a,\*</sup>, Lennart Bogg <sup>b</sup>, Clas Rehnberg <sup>c</sup>,  
Vinod Diwan <sup>b,d</sup>

Table 5  
Types of antibiotics by financing groups

Systems	Insurance (%)	Out-of-pocket (%)	Total (%)	$\chi^2$	<i>P</i>
Total cases				11.6	0.04
Penicillin	97 (37.9)	209 (45.3)	306 (42.7)		
Gentamicin	35 (13.7)	74 (16.1)	109 (15.2)		
Norfloxacin	27 (10.5)	37 (8.0)	64 (8.9)		
Cephalosporins	22 (8.6)	17 (3.7)	39 (5.4)		
Sulfonamides	13 (5.1)	21 (4.6)	34 (4.7)		
Others	62 (24.2)	103 (22.3)	165 (23.0)		
County hospital				2.0	0.16
Newer drugs	25 (30.5)	15 (20.5)	40 (25.8)		
Old drugs	57 (69.5)	58 (79.5)	115 (74.2)		
Township hospital				1.5	0.21
Newer drugs	16 (12.1)	30 (17.2)	46 (15.0)		
Old drugs	116 (87.9)	144 (82.8)	260 (85.0)		
Village health station					0.003*
Newer drugs	8 (19.0)	10 (4.7)	18 (7.0)		
Old drugs	34 (81.0)	204 (95.3)	238 (93.0)		
Respiratory infections				7.5	0.01
Newer drugs	20 (18.9)	19 (8.5)	39 (11.8)		
Old drugs	86 (81.1)	205 (91.5)	291 (88.2)		
Digestive system infections				1.0	0.32
Newer drugs	14 (23.0)	18 (16.7)	32 (18.9)		
Old drugs	47 (77.0)	90 (83.3)	137 (81.1)		

\* Fisher's exact test (two-tailed).

# Global availability of colistin



Wertheim et al, JGAR 2013



---

# Emergence of plasmid-mediated colistin resistance mechanism MCR-1 in animals and human beings in China: a microbiological and molecular biological study



Yi-Yun Liu\*, Yang Wang\*, Timothy R Walsh, Ling-Xian Yi, Rong Zhang, James Spencer, Yohei Doi, Guobao Tian, Baolei Dong, Xianhui Huang, Lin-Feng Yu, Danxia Gu, Hongwei Ren, Xiaojie Chen, Luchao Lv, Dandan He, Hongwei Zhou, Zisen Liang, Jian-Hua Liu, Jianzhong Shen


## Summary

**Background** Until now, polymyxin resistance has involved chromosomal mutations but has never been reported via horizontal gene transfer. During a routine surveillance project on antimicrobial resistance in commensal *Escherichia coli* from food animals in China, a major increase of colistin resistance was observed. When an *E coli* strain, SHP45, possessing colistin resistance that could be transferred to another strain, was isolated from a pig, we conducted further analysis of possible plasmid-mediated polymyxin resistance. Herein, we report the emergence of the first plasmid-mediated polymyxin resistance mechanism, MCR-1, in Enterobacteriaceae.

*Lancet Infect Dis* 2015

Published Online  
November 18, 2015  
[http://dx.doi.org/10.1016/S1473-3099\(15\)00424-7](http://dx.doi.org/10.1016/S1473-3099(15)00424-7)  
See Online/Articles  
[http://dx.doi.org/10.1016/S1473-3099\(15\)00462-6](http://dx.doi.org/10.1016/S1473-3099(15)00462-6)

## Countries reporting plasmid-mediated colistin resistance encoded by *mcr-1*



Isolate source(s):



Animals




Humans



Animals and humans



Animals and environment



Animals, humans  
and environment

Data source: Al-Tawfig, J. A., Lasminarayan, R. & Mendelson, M. How should we respond to the emergence of plasmid-mediated colistin resistance in humans and animals? *Int. J. Infect. Dis.* (2016). doi:10.1016/j.ijid.2016.11.415

III. Drivers of antibiotic use relate to incentives and behavior of patients, physicians, pharma, payers and healthcare institutions.

## Incentives for Physicians



- Satisfying patient expectations

TABLE 5

**Frequency of Antibiotic Prescribing by Factors Related to Patients' Expectations of Antibiotics (N = 482)**


Factor	No.* (%)	Antibiotic Prescribed No. (%)	OR (95% CI)
Patient expects antibiotic			
Yes	290 (60)	213 (73)	2.6 (1.7-3.9) reference
No	150 (31)	78 (52)	
No answer	42 (9)	28 (67)	
Clinician believes patient expects an antibiotic			
Yes	298 (62)	236 (79)	4.7 (3.2-7.1) reference
No	182 (38)	81 (45)	
No answer	2 (<1)	2 (100)	
Antibiotic helped similar illness in the past			
Yes	284 (59)	212 (75)	4.5 (2.9-6.9) reference
No	170 (35)	88 (52)	
Don't know	19 (4)	12 (63)	
No answer	9 (2)	5 (56)	

NOTE: Because some questions were unanswered, the numbers may not add up to 482.

\*In outpatients with nonspecific upper respiratory infections, acute bronchitis, or acute sinusitis.

OR denotes odds ratio; CI, confidence interval.

## Decision fatigue increases inappropriate prescribing




Relative to the first hour of a session, the adjusted odds ratios of antibiotic prescribing in the fourth hour was 1.26 (95% CI, 1.13–1.41)

Linder et al, JAMA IM, 2014

**Antibiotic restriction policy paradoxically increased private drug consumptions outside Taiwan's National Health Insurance**

Shu-Chen Kuo<sup>1,2</sup>, Shu-Men Shih<sup>3</sup>, Li-Yun Hsieh<sup>3</sup>,  
Tsai-Ling Yang<sup>4</sup>, Lauderdale<sup>5</sup>, Yee-Chun Chen<sup>1,2,3,4</sup>,  
Chao A. Hsiung<sup>2</sup> and Shou-Chwen Chong<sup>1,2</sup>



**Figure 1.** Changes in antibiotic consumption recorded in the IMS Health Taiwan database, relative to data from Taiwan's NHIRD. Biennial

## Hospital Incentives



- Antibiotics are a substitute for infection control
- Infection control is often not compensated



# Relationship Between Occurrence of Surgical Complications and Hospital Finances

---

Sunil Eappen, MD

---

Bennett H. Lane, MS

---

Barry Rosenberg, MD, MBA

---

Stuart A. Lipsitz, ScD

---

David Sadoff, MBA

---

Dave Matheson, JD, MBA

---

William R. Berry, MD, MPP, MPH

---

Mark Lester, MD, MBA

---

Atul A. Gawande, MD, MPH

---

**Importance** The effect of surgical complications on hospital finances is unclear.

**Objective** To determine the relationship between major surgical complications and per-encounter hospital costs and revenues by payer type.

**Design, Setting, and Participants** Retrospective analysis of administrative data for all inpatient surgical discharges during 2010 from a nonprofit 12-hospital system in the southern United States. Discharges were categorized by principal procedure and occurrence of 1 or more postsurgical complications, using *International Classification of Diseases, Ninth Revision*, diagnosis and procedure codes. Nine common surgical procedures and 10 major complications across 4 payer types were analyzed. Hospital costs and revenue at discharge were obtained from hospital accounting systems and classified by payer type.

Compared with absence of complications, complications were associated with a **\$39 017** higher contribution margin per patient with private insurance (\$55 953 vs \$16 936) and a **\$1749** higher contribution margin per patient with Medicare (\$3629 vs \$1880).



# Assessment of empirical antibiotic therapy optimisation in six hospitals: an observational cohort study

*Nikolay P Braykov\*, Daniel J Morgan\*, Marin L Schweizer, Daniel Z Uslan, Theodoros Kelesidis, Scott A Weisenberg, Birgir Johannsson, Heather Young, Joseph Cantey, Arjun Srinivasan, Eli Perencevich, Edward Septimus, Ramanan Laxminarayan*

## Summary

**Background** Modification of empirical antimicrobials when warranted by culture results or clinical signs is recommended to control antimicrobial overuse and resistance. We aimed to assess the frequency with which patients were started on empirical antimicrobials, characteristics of the empirical regimen and the clinical characteristics of patients at the time of starting antimicrobials. We also assessed the frequency of empirical therapy at different

*Lancet Infect Dis* 2014;  
14: 1220-27


See [Comment](#) page 1168


[Contributors](#)

- At the start of therapy, 220 (30%) patients were afebrile and had normal white blood cell counts.
- Appropriate cultures were collected from 432 (59%) patients, and 250 (58%) were negative.
- By the 5th day of therapy, 12.5% of empirical antimicrobials were escalated, 21.5% were narrowed or discontinued, and 66.4% were unchanged.
- Narrowing or discontinuation was more likely when cultures were collected at the start of therapy and no infection was noted on an initial radiological study.

IV. Antibiotic use in animal sector is increasing globally in response to the tremendous growth in demand for animal protein.


Meanwhile antibiotic manufacturing is expediting the accumulation of resistance genes in the environment.



A close-up photograph of a brown chicken with a prominent red comb, pecking at feed in a wire cage. The chicken is the central focus, with its head and neck in sharp detail. The background shows other chickens in similar cages, slightly out of focus. A semi-transparent grey box with a thin black border is overlaid on the left side of the image, containing text.

$2/3^{\text{rds}}$  of the  
tonnage of  
antibiotics sold  
worldwide are  
used in  
agriculture


Demand for poultry in India and China is set to increase two to seven fold between 2000 and 2030



# Pig Run


Swine output has surged to feed pork-hungry China

Per-capita Pork Consumption



Source: Bloomberg data

Number of Pigs Produced



Bloomberg 



## Drug Binge


China consumes half the world's antibiotics, with the majority administered to animals



Antibiotics consumed (metric tons) in 2013

Source: Ying Guang-Guo et al in Environmental Science & Technology, May 2015

Bloomberg 



In 2013, the global consumption of all antimicrobials in food animals was estimated at 131,109 tons

Projected to reach 200,235 tons by 2030

50,000 Tons

10,000 Tons

2,000 Tons

500 Tons


2030

2013

Van Boeckel et al, *Science* 2017

## Global antibiotic use in food animals by 2030


Capping the use of antibiotics at 50 milligrams (mg) per kilogram (kg) of animal product, reducing the consumption of meat to 165 grams (g) per day per person, and imposing a 50% tax on antibiotics could help reduce the use of antibiotics in food animals.



\*OECD—Organisation for Economic Co-operation and Development

Giorgia Guglielmi/Science; Data: T. P. Van Boeckel et al., Reducing antimicrobial use in food animals

Van Boeckel et al, *Science* 2017



- Total consumption in China - 92700 tons in 2013,
- 54000 tons of antibiotics excreted by human and animals - much of this entered into the receiving environment following various wastewater treatments into 58 river basins of China

Zhang et al, Env Sci Tech, 2015


# Diverse and abundant antibiotic resistance genes in Chinese swine farms

Yong-Guan Zhu<sup>a,b,1,2</sup>, Timothy A. Johnson<sup>c,d,1</sup>, Jian-Qiang Su<sup>a</sup>, Min Syed A. Hashsham<sup>c,e</sup>, and James M. Tiedje<sup>c,d,2</sup>

<sup>a</sup>Key Lab of Urban Environment and Health, Institute of Urban Environment, Chinese Academy of Sciences, Beijing 100085, China; and <sup>c</sup>Center for Environmental and Estuarine Science, and <sup>d</sup>Civil and Environmental Engineering, Michigan State University, East Lansing, Michigan 48824, USA

Contributed by James M. Tiedje, December 31, 2012 (sent for review October 31, 2012)


High-capacity quantitative PCR arrays detected 149 unique resistance genes among all of the farm samples, the top 63 ARGs being enriched 192-fold (median) up to 28,000-fold (maximum) compared with their respective antibiotic-free manure or soil controls.



# The Prevalence of Extended-Spectrum Beta-Lactamase-Producing Multidrug-Resistant *Escherichia Coli* in Poultry Chickens and Variation According to Farming Practices in Punjab, India

Charles H. Brower,<sup>1</sup> Siddhartha Mandal,<sup>2</sup> Shivdeep Hayer,<sup>3</sup> Mandeep Sran,<sup>4</sup> Asima Zehra,<sup>4</sup> Sunny J. Patel,<sup>5</sup> Ravneet Kaur,<sup>6</sup> Leena Chatterjee,<sup>7</sup> Savita Mishra,<sup>6</sup> B.R. Das,<sup>7</sup> Parminder Singh,<sup>8</sup> Randhir Singh,<sup>4</sup> J.P.S. Gill,<sup>4</sup> and Ramanan Laxminarayan<sup>1,9</sup>

Brower et al, *Environmental Health Perspectives*, 2017



## *Pharmaceuticals and Personal Care Products in the Environment*

### CONTAMINATION OF SURFACE, GROUND, AND DRINKING WATER FROM PHARMACEUTICAL PRODUCTION

JERKER FICK,\*† HANNA SÖDERSTRÖM,† RICHARD H. LINDBERG,† CHAU PHAN,† MATS TYSKLIND,† and D.G. JOAKIM LARSSON‡

†Department of Chemistry, Umeå University, Linneausväg 6, SE-90187 Umeå, Sweden

High amounts of four antibiotics were measured in the lakes that do not take in wastewater from the sewage plant. The levels of ciprofloxacin (2.5 mg/L) and cetirizine (20 µg/L) in one of the lakes was higher than previously measured levels in the blood of people taking the medications, report the authors. This suggests there are other unknown sources – perhaps illegal dumping – of wastewater responsible for polluting the lakes.

In addition, effluents from a wastewater treatment had concentrations of ciprofloxacin of 14 milligrams per liter (mg/L) and cetirizine as high as 1.2 mg/L. These concentrations are approaching therapeutic doses (concentrations that would kill some microorganisms outright). Concentration reported in the US range in the nanograms per liter (ng/L), which are one million fold less.


contaminated by the treatment plant. Water samples were also taken from wells in six nearby villages. The samples were analyzed for the presence of 12 pharmaceuticals with liquid chromatography–mass spectrometry. All wells were determined to be contaminated with drugs. Ciprofloxacin, enoxacin, cetirizine, terbinafine, and citalopram were detected at more than 1 µg/L in several wells. Very high concentrations of ciprofloxacin (14 mg/L) and cetirizine (2.1 mg/L) were found in the effluent of the treatment plant, together with high concentrations of seven additional pharmaceuticals. **Very high concentrations of ciprofloxacin (up to 6.5 mg/L), cetirizine (up to 1.2 mg/L), norfloxacin (up to 0.52 mg/L), and enoxacin (up to 0.16 mg/L) were also detected in the two lakes,** which clearly shows that the investigated area has additional environmental sources of insufficiently treated industrial waste. Thus, insufficient wastewater management in one of the world's largest centers for bulk drug production leads to unprecedented drug contamination of surface, ground, and drinking water. This raises serious concerns regarding the development of antibiotic resistance, and it creates a major challenge for producers and regulatory agencies to improve the situation.

Fick et al *Env Tox and Chem*, 2009

VICE  
NEWS









Increase of antibiotic resistance genes among soils collected at five sites in The Netherlands from 1940 to 2008.

## Antibiotic consumption in livestock, top ten countries 2010–2030 (projected for 2030)




Van Boeckel et al., PNAS, 2015



## Productivity reductions and costs per produced pig incurred by removing AGPs



V. Is finding new antibiotics the answer?

A close-up photograph of a person's hand holding a black gas pump nozzle. The nozzle is inserted into the fuel tank of a white car. The background is blurred, showing a red taillight and some greenery. The text "Make better use of existing antibiotics" is overlaid on the image in a blue-bordered box.

**Make better use of existing antibiotics**


Image Courtesy of Shutterstock

A photograph of an offshore oil rig at sunset. The rig's complex steel structure is silhouetted against a sky with soft orange and blue hues. The rig is supported by several legs in the water, and its reflection is visible on the calm surface. In the background, another smaller rig is visible on the horizon.

Find new antibiotics

Image Courtesy of Shutterstock


## Loss of first line drugs increases drug costs



Source: WHO Policy Perspective 2005, adapted from WHO Model Formulary, WHO Clinical Guidelines and Management Sciences for Health's 2004 International Drug Price Indicator Guide (slide courtesy: David Heymann)




# The rich pay with their wallets, the poor with their lives



Notes: \*Chloramphenicol is not available in developed world—price is therefore estimated. †Ceftriaxone and ciprofloxacin may be available in some tertiary settings in developing world.  
 Source: The Medical Letter (2006), Disease control priorities in developing countries, Lancet (2006), Expert interviews.

# Discovery of new classes of antibiotics




# Pipeline of new anti-microbial drugs growing after a long lag

But prices are likely to be high


Significant gap in first introduction of new antimicrobial classes...

Year	Class of drug
1935	Sulfonamides
1941	Penicillins
1944	Aminoglycosides
1945	Cephalosporins
1949	Chloramphenicol
1950	Tetracyclines
1952	Macrolides / Lincosamides / Streptogramins
1956	Glycopeptides
1957	Rifamycins
1959	Nitroimidazoles
1962	Quinolones
1968	Trimethoprim
<b>30 year development gap</b>	
2000	Oxazolidinones
2003	Lipopeptides
2005	Glycylcyclines
2007	Pleuromutilins

...Is being addressed by large recent reinvestment in novel mechanism development




# of molecules



# of molecules in non-standard anti-microbial classes in all phases of development


Source: "Bad Bugs, No Drugs" white paper, Pharmaprojects, Rodman and Renshaw, Nature Reviews Drug Discovery, BCG Analysis

### Total # New FDA Approved Antibacterial Agents, 1983-2014



Data sources: [idsociety.org](http://idsociety.org), FDA, *Nature Reviews* annual FDA drug approvals, 2006-2014

# Trends in development of new antibiotics



## Trends in development of new antibiotics



### New antibiotic launches since 1994

Launch Year	Product name	Antimicrobial class (old)	Pharmaceutical Company
		Antimicrobial class (new)	
1994	Meropenem	Carbapenem	AstraZeneca
1999	Moxifloxacin	Fluoroquinolone	Bayer
2000	Linezolid	Oxazolidinone	Pfizer
2001	Telithromycin	Macrolide	Sanofi-Aventis
2002	Balofloxacin	Fluoroquinolone	Choongwae Pharma
	Biapenem	Carbapenem	Wyeth
	Ertapenem	Carbapenem	Merck
	Prulifloxacin	Fluoroquinolone	Nippon Shinyaku Co.
	Pazufloxacin	Fluoroquinolone	Toyama Chemical Co.
2004	Gemifloxacin	Fluoroquinolone	LG Life Sciences
2005	Tigecycline	Glycylcycline	Wyeth
	Doripenem	Carbapenem	Janssen Pharmaceuticals
2006	Daptomycin	Lipopeptide	Cubist Pharmaceuticals
2007	Garenoxacin	Quinolone	Toyama Chemical Co.
	Retaparmulin	Pleuromutilin	GlaxoSmithKline
2008	Dalbavancin	Glycolipopeptide	Pfizer
	Oritavancin	Glycopeptide	Targanta Therapeutics
	Sitafloxacin	Fluoroquinolone	Daiichi Pharmaceutical Co.
	Telavancin	Novel glycolipopeptide	Theravance
2009	Antofloxacin	Fluoroquinolone	Anhui Global
	Besifloxacin	Fluoroquinolone	SSP Co.
	Ceftobiprole	5th-gen cephalosporin	Johnson & Johnson
	Iclaprim	DHFR inhibitor	Arpida
	Tebipenem	Carbapenem	Meiji Seika Pharma Co.
2011	Ceftaroline	5th-gen cephalosporin	Cerexa
	Fidaxomicin	Macrocyclic	Optimer Pharmaceuticals
2012	Bedaquiline	Diarylquinoline	Janssen Pharmaceuticals

Fig. 3. Antibiotic pipeline for the past 20 years.

Laxminarayan, *Science*, 2014

### New antibiotic launches since 1994

Launch Year	Product name	Antimicrobial class (old)	Antimicrobial class (new)	Pharmaceutical Company
1994	Meropenem	Carbapenem		AstraZeneca
1999	Moxifloxacin	Fluoroquinolone		Bayer
2000	Linezolid	Oxazolidinone		Pfizer
2001	Gemifloxacin	Fluoroquinolone		Wyeth
2002	Garenoxacin	Quinolone		Toyama Chemical Co.
2003	Retaparmulin	Pleuromutilin		GlaxoSmithKline
2004	Dalbavancin	Glycolipopeptide		Pfizer
2004	Gemifloxacin	Fluoroquinolone		LG Life Sciences
2005	Tigecycline	Glycylcycline		Wyeth
2005	Doripenem	Carbapenem		Janssen Pharmaceuticals
2006	Daptomycin	Lipopeptide		Cubist Pharmaceuticals
2007	Iclaprim	DHFR inhibitor		Arpida
2007	Tebipenem	Carbapenem		Meiji Seika Pharma Co.
2008	Ceftaroline	5th-gen cephalosporin		Cerexa
2009	Fidaxomicin	Macrocyclic		Optimer Pharmaceuticals
2011	Ceftaroline	5th-gen cephalosporin		Cerexa
2011	Fidaxomicin	Macrocyclic		Optimer Pharmaceuticals
2012	Bedaquiline	Diarylquinoline		Janssen Pharmaceuticals

Incentives for new antibiotics, as proposed by BARDA and EU may encourage new drug development but don't impact incentives for using drugs appropriately

Fig. 3. Antibiotic pipeline for the past 20 years.

Laxminarayan, *Science*, 2014



## BLADE OF GRASS IS RESPONSIBLE FOR LOSS OF FOOT

C. W. Jones, athletic director of the Athens Y. M. C. A. yesterday suffered the loss of his right foot, the member having been amputated just above the ankle.

Mr. Jones, it seems, recently was exercising on a plot of grass, dew on a blade of grass cutting him slightly just under the little toe. The cut did not heal as quickly as it should have and medical attention was called, but to no avail. Blood poisoning had set in, and it was imperative that the foot be amputated to prevent the poison spreading further.

*Weekly Banner*, 18 July 1899, p. 2, col. 2.  
©Athens-Clarke County Heritage Room, 2011.




Image Courtesy of Shutterstock





- Bacteriophages
- Probiotics
- Quorum sensing
- Synbiotics

**Price in USD**

\$ 40,000

\$ 20,000


\$ 200

\$ 100

\$ 0.20

\$ 0.10

0



Penicillin

**Market Launch: 1941**

**Price in USD**

\$ 40,000

\$ 20,000


\$ 200

\$ 100


\$ 0.20

\$ 0.10

0



Penicillin



Linezolid

**Market Launch: 1941**

**2000**

**Price in USD**

\$ 40,000

\$ 20,000


\$ 200

\$ 100

\$ 0.20

\$ 0.10

0



Penicillin

Linezolid


Daptomycin

**Market Launch: 1941**

**2000**

**2006**

**Price in USD**



**Market Launch: 1941**

**2000**

**2006**

**2010**

United Nations

A/71/L.2



## General Assembly

Distr.: Limited  
22 September 2016

Original: English

---

Seventy-first session

Agenda item 127


Global health and foreign policy


**Draft resolution submitted by the President of the General Assembly**

**Political Declaration of the high-level meeting of the General Assembly  
on antimicrobial resistance**



# Resistancemap.org





**Global Antibiotic Resistance  
Partnership (GARP) Member Countries**

Slides are @ [www.cddep.org](http://www.cddep.org)

Thank you